Information for breastfeeding families

Plugged Ducts & Mastitis



Plugged Ducts

If you notice a small lump the size of a pea in your breasts, it may be a plugged duct. This occurs when a portion of the breast does not get emptied completely during feedings.

Remedy:

- ✓ Apply a warm compress to the area before feeding
- ✓ Massage the lump towards the nipple during a feeding
- ✓ It may take 2 or 3 feedings for it to completely empty. Position your baby's chin towards the area of the lump. This is where the greatest emptying will occur.
- ✓ If you find a persistent lump that does not respond to these measures, please see your physician. It could be a different problem.

Plugged Nipple Pore

This appears as a small white dot on the tip of the nipple and is usually very painful. It is one milk duct that has become plugged.



Remedy:

- ✓ Soften the plug with a warm compress
- ✓ Massage the nipple beginning near the plug and gradually work your way back following the duct, if you can feel it.
- ✓ In persistent cases, soak the nipple with olive oil on a cotton ball for several hours, then massage the nipple and gently scrape the skin surface while in the shower

Mastitis

This occurs most frequently in mothers who have had a cracked or blistered nipple or who are undergoing a period of stress such as returning to work, participating in holiday activities or experiencing a change in normal daily routine.

Symptoms may include:

- High fever, starting suddenly Hot, reddened area
- Red streaks
- Pain and a lump in the breast Hard wedge shaped area Flu like symptoms chills Extreme tiredness





Remedy

- ✓ Apply warm compresses before feedings
- ✓ Gently massage the area
- √ Keep your breast empty by frequent nursing
- ✓ Continue to breastfeed, even on the affected side
- ✓ If your baby does not empty that side well, use a good quality breast pump after feedings
- ✓ Apply ice after feedings for 10 15 minutes for the first day or two.
- ✓ Rest in bed as much as you can
- ✓ Drink plenty of fluids.

Your physician, usually your obstetrician or family doctor, will prescribe an antibiotic. You must take a full 7 -10 day course of medication. Do not stop taking it until the prescription is gone even though you start to feel better. Inadequately treated mastitis is more likely to return.

Provided for you by Lactation Education Resources. This handout may be freely duplicated. Please be aware that the information provided is intended solely for general educational and informational purposes only. It is neither intended nor implied to be a substitute for professional medical advice. Always seek the advice of your physician for any questions you may have regarding you or your infant's medical condition. Never disregard professional medical advice or delay in seeking it because of something you have received in this information.